

Hindawi Publishing Corporation  
Advances in Difference Equations  
Volume 2011, Article ID 913403, 6 pages  
doi:10.1155/2011/913403

## Research Article

# Notes on Interpolation Inequalities

Jiu-Gang Dong<sup>1</sup> and Ti-Jun Xiao<sup>2</sup>

<sup>1</sup> Department of Mathematics, University of Science and Technology of China, Hefei 230026, China

<sup>2</sup> Shanghai Key Laboratory for Contemporary Applied Mathematics, School of Mathematical Sciences, Fudan University, Shanghai 200433, China

Correspondence should be addressed to Ti-Jun Xiao, [xiaotj@ustc.edu.cn](mailto:xiaotj@ustc.edu.cn)

Received 3 October 2010; Accepted 16 November 2010

Academic Editor: Toka Diagana

Copyright © 2011 J.-G. Dong and T.-J. Xiao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

An easy proof of the John-Nirenberg inequality is provided by merely using the Calderón-Zygmund decomposition. Moreover, an interpolation inequality is presented with the help of the John-Nirenberg inequality.

## 1. Introduction

It is well known that various interpolation inequalities play an important role in the study of operational equations, partial differential equations, and variation problems (see, e.g., [1–6]). So, it is an issue worthy of deep investigation.

Let  $Q_0$  be either  $R^n$  or a fixed cube in  $R^n$ . For  $f \in L^1_{\text{loc}}(Q_0)$ , write

$$\|f\|_{\text{BMO}} := \sup_{Q \subset Q_0} \frac{1}{|Q|} \int_Q |f - f_Q| dx, \quad (1.1)$$

where the supremum is taken over all cubes  $Q \subset Q_0$  and  $f_Q := (1/|Q|) \int_Q f dx$ .

Recall that  $\text{BMO}(Q_0)$  is the set consisting of all locally integrable functions on  $Q_0$  such that  $\|f\|_{\text{BMO}} < \infty$ , which is a Banach space endowed with the norm  $\|\cdot\|_{\text{BMO}}$ . It is clear that any bounded function on  $Q_0$  is in  $\text{BMO}(Q_0)$ , but the converse is not true. On the other hand, the BMO space is regarded as a natural substitute for  $L^\infty$  in many studies. One of the important features of the space is the John-Nirenberg inequality. There are several versions of its proof; see, for example, [2, 7–9]. Stimulated by these works, we give, in this paper, an easy proof of the John-Nirenberg inequality by using the Calderón-Zygmund decomposition only. Moreover, with the help of this inequality, an interpolation inequality is showed for  $L^p$  and BMO norms.

## 2. Results and Proofs

**Lemma 2.1** (John-Nirenberg inequality). *If  $f \in BMO(Q_0)$ , then there exist positive constants  $c_1, c_2$  such that, for each cube  $Q \subset Q_0$ ,*

$$|\{x \in Q : |f(x) - f_Q| > t\}| \leq c_1 \exp\left\{-\frac{c_2}{\|f\|_{BMO}} t\right\} |Q|, \quad t > 0. \quad (2.1)$$

*Proof.* Without loss of generality, we can and do assume that  $\|f\|_{BMO} = 1$ .

For each  $t > 0$ , let  $F(t)$  denote the least number for which we have

$$|\{x \in Q : |f(x) - f_Q| > t\}| \leq F(t)|Q|, \quad (2.2)$$

for any cube  $Q \subset Q_0$ . It is easy to see that  $F(t) \leq 1$  ( $t > 0$ ) and  $F(t)$  is decreasing.

Fix a cube  $Q \subset Q_0$ . Applying the Calderón-Zygmund decomposition (cf., e.g., [2, 9]) to  $|f(x) - f_Q|$  on  $Q$ , with  $2^n$  as the separating number, we get a sequence of disjoint cubes  $\{Q_j\}$  and  $E$  such that

$$Q = \left(\bigcup_j Q_j\right) \cup E, \quad (2.3)$$

$$|f(x) - f_Q| \leq 2^n, \quad \text{for a.e. } x \in E, \quad (2.4)$$

$$2^n < \frac{1}{|Q_j|} \int_{Q_j} |f - f_Q| dx \leq 4^n. \quad (2.5)$$

Using (2.5), we have

$$\sum_j |Q_j| < \frac{1}{2^n} |Q|. \quad (2.6)$$

From (2.3), (2.4), and (2.6), we deduce that for  $t > 4^n$ ,

$$\begin{aligned} |\{x \in Q : |f(x) - f_Q| > t\}| &= \left| \bigcup_j \{x \in Q_j : |f(x) - f_Q| > t\} \right| \\ &\leq \sum_j |\{x \in Q_j : |f(x) - f_Q| > t - 4^n\}| \\ &= \sum_j |Q_j| \frac{1}{|Q_j|} |\{x \in Q_j : |f(x) - f_Q| > t - 4^n\}| \\ &\leq \frac{1}{2^n} F(t - 4^n) |Q|. \end{aligned} \quad (2.7)$$

This yields that

$$F(t) \leq \frac{1}{2^n} F(t - 4^n), \quad t > 4^n. \quad (2.8)$$

Let  $\gamma = [(t-1)4^{-n}]$  ( $t > 4^n$ ),  $\mu = 1 + \gamma 4^n$ . Then  $0 < \mu \leq t$ . By iterating, we get

$$\begin{aligned} F(t) &\leq F(\mu) = F(1 + \gamma 4^n) \leq 2^{-n\gamma} \leq 2^{-n((t-1)4^{-n}-1)} \\ &= 2^{n(1+4^{-n})} \exp(-(\log 2)n4^{-n}t), \quad t > 4^n. \end{aligned} \quad (2.9)$$

Thus, letting

$$c_1 = 2^{n(1+4^{-n})}, \quad c_2 = (\log 2)n4^{-n} \quad (2.10)$$

gives that

$$F(t) \leq c_1 e^{-c_2 t}, \quad t > 0, \quad (2.11)$$

since

$$F(t) \leq 1 \leq c_1 e^{-c_2 t}, \quad 0 < t \leq 4^n. \quad (2.12)$$

This completes the proof.  $\square$

*Remark 2.2.* (1) As we have seen, the recursive estimation (2.8) justifies the desired exponential decay of  $F(t)$ .

(2) There exists a gap in the proof of the John-Nirenberg inequality given in [2]. Actually, for a decreasing function  $G(t) : (0, \infty) \rightarrow [0, 1]$ , the following estimate:

$$G(2 \cdot 2^n \alpha) \leq \frac{1}{\alpha} G(2^n \alpha), \quad \alpha > 1 \quad (2.13)$$

does not generally imply such a property, that is, the existence of constants  $c_1, c_2 > 0$  such that

$$G(t) \leq c_1 e^{-c_2 t}, \quad t > 0. \quad (2.14)$$

We present the following function as a counter example:

$$G(t) = \exp \left\{ - \left( \log \frac{5}{3} \right)^{-1} \log^2(t+1) \right\}, \quad t > 0. \quad (2.15)$$

In fact, it is easy to see that there are no constants  $c_1, c_2 > 0$  such that (2.14) holds. On the other hand, we have

$$\frac{G'(t)}{G(t)} = \left\{ -\left(\log \frac{5}{3}\right)^{-1} 2 \frac{\log(t+1)}{t+1} \right\}, \quad t > 0. \quad (2.16)$$

Integrating both sides of the above equation from  $2^n \alpha$  to  $2 \cdot 2^n \alpha$ , we obtain

$$\begin{aligned} G(2 \cdot 2^n \alpha) &= \exp \left\{ -2 \left(\log \frac{5}{3}\right)^{-1} \int_{2^n \alpha}^{2 \cdot 2^n \alpha} \frac{\log(t+1)}{t+1} dt \right\} G(2^n \alpha) \\ &= \exp \left\{ -\left(\log \frac{5}{3}\right)^{-1} \left( \log^2(2 \cdot 2^n \alpha + 1) - \log^2(2^n \alpha + 1) \right) \right\} G(2^n \alpha) \\ &= \exp \left\{ -\left(\log \frac{5}{3}\right)^{-1} \log((2 \cdot 2^n \alpha + 1)(2^n \alpha + 1)) \cdot \log\left(\frac{2 \cdot 2^n \alpha + 1}{2^n \alpha + 1}\right) \right\} G(2^n \alpha) \quad (2.17) \\ &\leq \exp \{ -\log((2 \cdot 2^n \alpha + 1)(2^n \alpha + 1)) \} G(2^n \alpha) \\ &= \frac{1}{(2 \cdot 2^n \alpha + 1)(2^n \alpha + 1)} G(2^n \alpha) \\ &\leq \frac{1}{\alpha} G(2^n \alpha), \end{aligned}$$

where the fact that

$$\frac{2 \cdot 2^n \alpha + 1}{2^n \alpha + 1} > \frac{5}{3} \quad (\alpha > 1) \quad (2.18)$$

is used to get the first inequality above. This means that

$$G(2 \cdot 2^n \alpha) \leq \frac{1}{\alpha} G(2^n \alpha), \quad \alpha > 1. \quad (2.19)$$

Next, we make use of the John-Nirenberg inequality to obtain an interpolation inequality for  $L^p$  and BMO norms.

**Theorem 2.3.** *Suppose that  $1 \leq p < r < \infty$  and  $f \in L^p(Q_0) \cap BMO(Q_0)$ . Then we have*

$$\|f\|_{L^r} \leq (\text{const}) \|f\|_{L^p}^{p/r} \|f\|_{BMO}^{1-p/r}. \quad (2.20)$$

*Proof.* If  $\|f\|_{\text{BMO}} = 0$ , the proof is trivial; so we assume that  $\|f\|_{\text{BMO}} \neq 0$ . In view of the Calderón-Zygmund decomposition theorem, there exists a sequence of disjoint cubes  $\{Q_j\}$  and  $E$  such that

$$Q_0 = \left( \bigcup_j Q_j \right) \cup E, \quad (2.21)$$

$$|f(x)|^p \leq \|f\|_{\text{BMO}}^p \quad \text{for a.e. } x \in E, \quad (2.22)$$

$$\|f\|_{\text{BMO}}^p < \frac{1}{|Q_j|} \int_{Q_j} |f(x)|^p dx \leq 2^n \|f\|_{\text{BMO}}^p. \quad (2.23)$$

From (2.23), we get

$$\begin{aligned} \sum_j |Q_j| &< \frac{1}{\|f\|_{\text{BMO}}^p} \int_{Q_0} |f(x)|^p dx = \frac{\|f\|_{L^p}^p}{\|f\|_{\text{BMO}}^p}, \\ |f|_{Q_j} &= \frac{1}{|Q_j|} \int_{Q_j} |f(x)| dx \leq \left( \frac{1}{|Q_j|} \int_{Q_j} |f(x)|^p dx \right)^{1/p} \leq 2^{n/p} \|f\|_{\text{BMO}}. \end{aligned} \quad (2.24)$$

Using (2.21)–(2.24), together with Lemma 2.1, yields that, for  $\lambda > 2^{n/p} \|f\|_{\text{BMO}}$ ,

$$\begin{aligned} |\{x \in Q_0 : |f(x)| > \lambda\}| &= \left| \bigcup_j \{x \in Q_j : |f(x)| > \lambda\} \right| \\ &\leq \sum_j \left| \{x \in Q_j : |f(x) - f_{Q_j}| > \lambda - |f_{Q_j}|\} \right| \\ &\leq \sum_j |Q_j| \frac{1}{|Q_j|} \left| \{x \in Q_j : |f(x) - f_{Q_j}| > \lambda - 2^{n/p} \|f\|_{\text{BMO}}\} \right| \quad (2.25) \\ &\leq \sum_j c_1 \exp \left\{ -\frac{c_2}{\|f\|_{\text{BMO}}} (\lambda - 2^{n/p} \|f\|_{\text{BMO}}) \right\} |Q_j| \\ &\leq c_1 \exp \left\{ -\frac{c_2}{\|f\|_{\text{BMO}}} (\lambda - 2^{n/p} \|f\|_{\text{BMO}}) \right\} \frac{\|f\|_{L^p}^p}{\|f\|_{\text{BMO}}^p}. \end{aligned}$$

From (2.25), we obtain

$$\begin{aligned}
 \|f\|_{L^r}^r &= r \int_0^\infty \lambda^{r-1} |\{x \in Q_0 : |f(x)| > \lambda\}| d\lambda \\
 &= r \int_0^{2^{n/p}\|f\|_{\text{BMO}}} \lambda^{r-1} |\{x \in Q_0 : |f(x)| > \lambda\}| d\lambda \\
 &\quad + r \int_{2^{n/p}\|f\|_{\text{BMO}}}^\infty \lambda^{r-1} |\{x \in Q_0 : |f(x)| > \lambda\}| d\lambda \\
 &\leq r \int_0^{2^{n/p}\|f\|_{\text{BMO}}} \lambda^{r-1} \frac{\|f\|_{L^p}^p}{\lambda^p} d\lambda \\
 &\quad + r \int_{2^{n/p}\|f\|_{\text{BMO}}}^\infty \lambda^{r-1} c_1 \exp\left\{-\frac{c_2}{\|f\|_{\text{BMO}}}(\lambda - 2^{n/p}\|f\|_{\text{BMO}})\right\} \frac{f\|f\|_{L^p}^p}{\|f\|_{\text{BMO}}^p} d\lambda \\
 &= \frac{r}{r-p} 2^{(n/p)(r-p)} \|f\|_{\text{BMO}}^{r-p} \|f\|_{L^p}^p + \frac{rc_1}{c_2} 2^{(n/p)(r-1)} \|f\|_{\text{BMO}}^{r-p} \|f\|_{L^p}^p \\
 &\leq (\text{const}) \|f\|_{\text{BMO}}^{r-p} \|f\|_{L^p}^p.
 \end{aligned} \tag{2.26}$$

Hence, the proof is complete.  $\square$

## Acknowledgments

The authors would like to thank the referees for helpful comments and suggestions. The work was supported partly by the NSF of China (11071042) and the Research Fund for Shanghai Key Laboratory for Contemporary Applied Mathematics (08DZ2271900).

## References

- [1] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Classics in Mathematics, Springer, Berlin, Germany, 2001.
- [2] Y. S. Han, *Modern Harmonic Analysis Methods and Applications*, Science Press, Beijing, China, 1997.
- [3] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, vol. 18 of *North-Holland Mathematical Library*, North-Holland, Amsterdam, The Netherlands, 1978.
- [4] T.-J. Xiao and J. Liang, *The Cauchy Problem for Higher-Order Abstract Differential Equations*, vol. 1701 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1998.
- [5] T.-J. Xiao and J. Liang, "Complete second order differential equations in Banach spaces with dynamic boundary conditions," *Journal of Differential Equations*, vol. 200, no. 1, pp. 105–136, 2004.
- [6] T.-J. Xiao and J. Liang, "Second order parabolic equations in Banach spaces with dynamic boundary conditions," *Transactions of the American Mathematical Society*, vol. 356, no. 12, pp. 4787–4809, 2004.
- [7] C. George, *Exercises in Integration*, Springer, New York, NY, USA, 1984.
- [8] F. John and L. Nirenberg, "On functions of bounded mean oscillation," *Communications on Pure and Applied Mathematics*, vol. 14, pp. 415–426, 1961.
- [9] Elias M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, vol. 43 of *Princeton Mathematical Series*, Princeton University Press, Princeton, NJ, NY, USA, 1993.